

***Dietary Vitamin D3, the Skin Immune System and
implications for Skin Cancer***

by

Dr Roslyn C Malley, B. Med. Sci (Hons), MBBS (Hons)

Submitted in fulfilment of the requirements for the degree of Doctor of Philosophy.

University of Tasmania
June 2010

Declaration of Originality

This thesis contains no material that has been accepted for a degree or diploma by the University or any other institution, except by way of background information and duly acknowledged in the thesis, and to the best of my knowledge and belief no material previously published or written by another person except where due reference is made in the text of this thesis, nor does the thesis contain any material that infringes copyright.

Roslyn Malley

Authority of Access

This thesis may be made available for loan and limited copying in accordance with the Copyright Act 1968.

Roslyn Malley

Part of the work from this thesis has been published or presented at conferences as follows:

Publications:

Malley, R.C., Muller H.K., Norval, M., Woods, G.M. (2009) Vitamin D3 deficiency enhances contact hypersensitivity in male but not in female mice. *Cellular Immunol* 255: 33-40

Talks:

Malley, R.C., Muller H.K., Woods, G.M. 'Dietary Vitamin D3 protects against UVB-induced immunosuppression in C57BL/6, but not BALB/c mice'. 5th Joint Meeting of the Societies for Free Radical Australasia and Japan & mutagenesis and Experimental Pathology Society of Australasia, Sydney, Australia, 2009

Malley, R.C., Muller H.K., Woods, G.M. 'Vitamin D3 nutrition influences regulatory T cell induction in neonatal tolerance and adult immunity'. Australasian Medical Health and Research Conference, Brisbane, 2008

Malley, R.C., Muller H.K., Woods, G.M. 'Dietary Vitamin D3 Influences UVB Induced DNA Damage and Immunosuppression'. Mutagenesis and Experimental Pathology of Australasia Conference, Hobart, 2007

Malley, R.C., Muller H.K., Norval, M., Woods, G.M. 'Modification of Skin Immune System Function by Dietary Vitamin D3 is influenced by Age and Gender'. National Health and Medical Research Congress, Melbourne, 2006

Poster presentations:

Malley, R.C., Muller, H.K., Woods, G.M. 'Vitamin D Promotes Neonatal Tolerance Induction to Contact Sensitisers'. Presented at the Australasian Society for Immunology Meeting, Manly, 2007

Malley, R.C., Muller, H.K. and Woods, G.M. '25-hydroxyvitamin D3 deficiency affects the contact sensitivity response differentially, depending on age and gender'. Presented at the 13th Vitamin D Workshop, Victoria, Canada, April, 2006.

Abstract

Ultraviolet radiation (UVR) causes mutagenic DNA damage to skin cells and suppression of the skin immune system (SIS) that contributes to skin cancer development. UVR in the UVB range also causes the beneficial production of vitamin D3 by keratinocytes. Keratinocytes also contain the enzymes required for the two-step hydroxylation of vitamin D3 to its biologically active form: 1 α ,25 dihydroxyvitamin D3 (1 α ,25 (OH)₂D3). 1 α ,25 (OH)₂D3 has been demonstrated to modulate the SIS and its response to UVB irradiation. Topical application of 1 α ,25 (OH)₂D3 suppresses the SIS and protects against UV-induced DNA damage. This thesis investigates the hypothesis that dietary vitamin D3 similarly suppresses the SIS response via modulation of the SIS and protects against UVB induced immunosuppression via a reduction in UVB induced DNA damage.

To assess the influence of vitamin D3 on the development and function of the SIS, and the skins response to UVR, a vitamin D3 deficient mouse population was developed. Breeding pairs of vitamin D3 replete and vitamin D3 deficient mice were formed and the off-spring of these mice were used in experiments. The contact hypersensitivity (CHS) response was used to assess the SIS function in vitamin D3 replete and vitamin D3 deficient mice. In this model the application of the contact sensitizer to the skin of neonatal mice leads to suppression of the CHS response upon subsequent re-sensitisation and elicitation of the CHS in adulthood. The suppression of the CHS response is termed 'neonatal tolerance'.

Vitamin D3 enhanced neonatal tolerance in male but not female mice. The enhanced neonatal tolerance induction related to an increase in the proportion of CD4⁺CD25⁺ cells in vitamin D3 replete neonatal male mice and this cell population could transfer tolerance in this mouse group. Vitamin D3 suppressed the CHS response in adult male but not adult female mice. An investigation of DC function in adult mice revealed no modulation by vitamin D3 in antigen carriage or co-stimulatory molecule expression, suggesting that these cells were not modulating the reduced CHS response in adult male mice. However, dietary vitamin D3 was associated with a higher proportion of

CD4⁺CD25⁺ cells in the SDLN after contact sensitiser application in male but not female mice, implying that vitamin D3 was modulating the induction of these cells.

The level of UVB-induced immunosuppression was assessed in vitamin D3 replete and deficient mice in C57BL/6 and BALB/c mice. Vitamin D3 protected C57BL/6, but not BALB/c mice against UVB-induced immunosuppression. The protection against UVB-induced immunosuppression may relate to a reduction in UVB-induced DNA damage in C57BL/6 mice, not identified in BALB/c mice. In BALB/c mice female gender protected against UVB-induced immunosuppression and was associated with a subtle lowering of UVB-induced DNA damage in comparison to male mice at low UVR exposures. These results indicate that the protection against UVB-induced immunosuppression by vitamin D3 varies with genetic background and the modulation of UVB-induced DNA damage.

If these mouse studies are extended to humans, the ability of vitamin D3 to suppress the SIS and the limited protection against UVB-induced DNA damage in males, depending on their genetic background, may contribute to the greater burden of skin cancer in males than in females.

Acknowledgements

The work presented in this thesis would not be possible had it not been for the support, faith, good humour, patience and encouragement of my supervisors Associate Professor Greg Woods and Emeritus Professor Konrad Muller. Their hard work is immensely appreciated and I cannot express my gratitude enough. A special thanks to Professor Mary Norval for her kind words and feedback.

The development and maintenance of the vitamin D3 deficient mouse colony would not have been possible without the assistance of Specialty Feeds, Central Animal House Staff, Jill Chuckowree and the Royal Hobart Hospital Endocrinology Department. I am grateful for their care and attention of the mouse food, mice and vitamin D3 testing.

Thanks to Mark Cozens for his expertise in flow cytometry, humour and company during the course of many samples. Thanks to the Royal Hobart Hospital Anatomical Pathology Department for their support, expertise and use of equipment. Thanks also to Narelle Phillips for expertly sectioning my numerous tissue specimens with a smile.

Thankyou to the Royal Hobart Hospital Research Foundation, Cancer Council of Tasmania, Arthritis Foundation of Tasmania, the National Health and Research Council and the Royal College of Pathologists of Australasia for their financial assistance.

Special thanks to colleagues, family and friends that have contributed their time and humour, especially during the writing phase, which has allowed me to complete this work. To those colleagues that have been there from the start: Nick, Che, Deborah, Kylie and Alex - what a journey we have had - I can not thank you all enough.

Finally, a special thanks to Dr Heather McGee, my sister Bronwyn Malley and partner Alastair Lamont. Their kind and generous spirits, along with my supervisors, have assisted me in reaching this point.

Commonly Used Abbreviations

APC	Antigen Presenting Cells
BCC	Basal Cell Carcinoma
CHS	Contact Hypersensitivity
ConA	Concanavalin A
CCR	Chemokine Receptor
CCL	Chemokine Ligand
cDC	conventional or classical Dendritic Cell
CTLA-4	Cytotoxic T-lymphocyte Antigen 4
CLA	Cutaneous Lymphocyte Antigen
CPD	Cyclobutane Pyrimidine Dimer
DC	Dendritic Cells
7-DHC	7- dehydrocholesterol
EAE	Experimental Autoimmune Encephalomyelitis
FITC	Fluorescein Isothiocyanate
Flt3	fms-related tyrosine kinase 3
IBD	Inflammatory Bowel Disease
IFN	Interferon
IL	Interleukin
iNKT	invariant Natural Killer T cell
iTreg	induced regulatory T cell
LC	Langerhans cell
MCP-1	Monocyte Chemoattractant Protein- 1
MHC	Major Histocompatibility Complex
MMP	Matrix Metalloproteinase
MS	Multiple Sclerosis
MT	Metallothionein
NK	Natural Killer
NMSC	Non-Melanoma Skin Cancer
NOD	Non-obese Diabetic
nTregs:	naturally occurring regulatory T cells

PAF	Platelet Activating Factor
pDC	plasmacytoid Dendritic Cell
PGE2	Prostaglandin E2
RANK	Receptor Activator for Nuclear Factor κ B
RANKL	Receptor Activator for Nuclear Factor κ B Ligand
ROS	Reactive Oxygen Species
SDLN	Skin Draining Lymph Nodes
SEM	Standard Error of the Mean
SD	Standard Deviation
SCC	Squamous Cell Carcinoma
SIS	Skin Immune System
TD	Thymine Dimers
TCR	T Cell Receptor
Th	T helper
TGF- β	Transforming Growth Factor β
Treg	Regulatory T cell
UCA	Urocanic Acid
UV	Ultraviolet
UVR	Ultraviolet Radiation
VDR	Vitamin D Receptor

Table of Contents

FOREWORD	3
1 CHAPTER 1: LITERATURE REVIEW	4
1.1 INTRODUCTION.....	4
1.2 THE VITAMIN D ENDOCRINE SYSTEM.....	6
1.3 ACTIONS OF THE VITAMIN D3 ENDOCRINE SYSTEM ON THE DEVELOPMENT AND FUNCTION OF THE ADAPTIVE IMMUNE SYSTEM.....	16
1.4 ORGANIZATION OF THE SKIN IMMUNE SYSTEM (SIS)	23
1.5 UVB, VITAMIN D AND SKIN CANCER DEVELOPMENT	35
1.6 AIMS OF THE THESIS.....	48
2 CHAPTER 2: MATERIALS AND METHODS	51
2.1 REAGENTS, CONSUMABLES AND EQUIPMENT.....	51
2.2 SOLUTIONS	54
2.3 ANIMALS	59
2.4 TREATMENT OF MICE	60
2.5 CELL TRANSFER EXPERIMENTS.....	63
2.6 EPIDERMAL SHEETS.....	64
2.7 CELL PREPARATIONS.....	65
2.8 MAGNETIC CELL SEPARATION	66
2.9 ANTIBODY LABELLING FOR FLOW CYTOMETRY	68
2.10 FUNCTIONAL ASSAYS	71
2.11 CYTOKINE ANALYSIS	72
2.12 TISSUE PROCESSING FOR HISTOLOGY AND IMMUNOHISTOCHEMISTRY.....	74
2.13 ANALYSIS OF TISSUE SECTIONS	77
2.14 STATISTICAL ANALYSIS	78
3 CHAPTER 3: CHARACTERISATION OF THE VITAMIN D3 DEFICIENT POPULATION AND THE CONTACT HYPERSENSITIVITY RESPONSE	79
3.1 INTRODUCTION.....	79
3.2 CHARACTERISTICS OF THE D+ AND D- BREEDING COLONY	80
3.3 EXPERIMENTAL ADULT D+ AND ADULT D- MOUSE 25(OH)D AND CALCIUM LEVELS	81
3.4 MORPHOLOGICAL CHARACTERISTICS OF NEONATAL AND ADULT EXPERIMENTAL MICE.....	82
3.5 CONTACT HYPERSENSITIVITY RESPONSE AND NEONATAL TOLERANCE INDUCTION	86
3.6 HISTOLOGY OF NON-SPECIFIC INFLAMMATION IN ADULT MICE.....	89
3.7 HISTOLOGY OF THE CONTACT HYPERSENSITIVITY RESPONSE IN IMMUNE ADULT AND NEONATALLY TOLERISED MICE.....	90
3.8 DISCUSSION.....	93
4 CHAPTER 4: INFLUENCE OF VITAMIN D3 ON DENDRITIC CELL DEVELOPMENT AND FUNCTION IN THE SKIN IMMUNE SYSTEM.....	102
4.1 INTRODUCTION.....	102
4.2 CHARACTERISATION OF THE DEVELOPMENT OF THE EPIDERMAL LANGERHANS CELL NETWORK	103
4.3 CHARACTERISATION OF THE SKIN DRAINING LYMPH NODE DENDRITIC CELLS IN NAÏVE MICE..	104
4.4 CO-STIMULATORY MOLECULE EXPRESSION BY THE SDLN DC SUBSETS IN NAÏVE MICE.....	108
4.5 DENDRITIC CELL MIGRATION DURING THE SENSITISATION PHASE OF THE CHS RESPONSE TO OXAZOLONE IN ADULT MICE	110
4.6 THE CHARACTERISATION OF SKIN DRAINING LYMPH NODE DENDRITIC CELLS 5 DAYS AFTER THE APPLICATION OF OXAZOLONE	111
4.7 CO-STIMULATORY EXPRESSION OF THE SDLN DC SUBSETS IN NEONATAL AND ADULT MICE 5 DAYS AFTER THE APPLICATION OF OXAZOLONE.....	115
4.8 DENDRITIC CELL FUNCTION, ANTIGEN CARRIAGE/ACQUISITION AND CO-STIMULATORY MOLECULE EXPRESSION IN ADULT MICE.....	116
4.9 DISCUSSION	118
5 CHAPTER 5: INFLUENCE OF VITAMIN D3 ON LYMPHOCYTE DEVELOPMENT AND FUNCTION	127
5.1 INTRODUCTION.....	127

5.2	LYMPHOID CELLS WITHIN THE SKIN DRAINING LYMPH NODES OF NAÏVE NEONATAL AND ADULT MICE.....	128
5.3	CONA STIMULATED CELL PROLIFERATION OF NAÏVE NEONATAL AND ADULT D+ AND D-, MALE AND FEMALE MICE LYMPH NODE OR SPLEEN CELL SUSPENSIONS	134
5.4	CYTOKINE PRODUCTION OF CONA STIMULATED SPLEEN AND LYMPH NODE SUSPENSIONS	135
5.5	DISCUSSION.....	142
6	CHAPTER 6: INFLUENCE OF VITAMIN D3 ON LYMPHOCYTE POPULATIONS IN THE CONTACT HYPERSENSITIVITY RESPONSE.....	150
6.1	INTRODUCTION.....	150
6.2	CHARACTERISATION OF LYMPHOCYTES FROM THE SKIN DRAINING LYMPH NODE 18 HOURS AFTER THE APPLICATION OF VEHICLE OR OXAZOLONE	151
6.3	CHARACTERISATION OF PROLIFERATION AND IFN- γ PRODUCTION BY SKIN DRAINING LYMPH NODE CELLS 18 HOURS AFTER THE APPLICATION OF VEHICLE OR OXAZOLONE	155
6.4	CHARACTERISATION OF LYMPHOCYTES FROM THE SKIN DRAINING LYMPH NODES 5 DAYS AFTER THE APPLICATION OF OXAZOLONE	156
6.5	CHARACTERISATION OF CONCAVALIN A-STIMULATED PROLIFERATION OF SKIN DRAINING LYMPH NODE AND SPLEEN CELLS FROM ADULT MICE 5 DAYS AFTER THE APPLICATION OF OXAZOLONE.....	168
6.6	CHARACTERISATION OF THE CYTOKINE PRODUCTION BY CONCAVALIN A STIMULATED SKIN DRAINING LYMPH NODE AND SPLEEN CELL SUSPENSIONS 5 DAYS AFTER THE APPLICATION OF OXAZOLONE.....	171
6.7	FOXP3 ⁺ CELL INFILTRATE IN THE EARS OF NEONATALLY TOLERISED AND IMMUNE MICE.....	178
6.8	TRANSFER OF NEONATAL TOLERANCE	179
6.9	DISCUSSION.....	180
7	CHAPTER 7: VITAMIN D3 AND THE SKINS RESPONSE TO UVB IRRADIATION	192
7.1	INTRODUCTION.....	192
7.2	UVB-INDUCED IMMUNOSUPPRESSION IN BALB/C AND C57BL/6 MICE.....	193
7.3	UVB-INDUCED DNA DAMAGE.....	195
7.4	REPAIR OF UVB-INDUCED DNA DAMAGE.....	198
7.5	VITAMIN D3 AND SKIN HYPERPLASIA AFTER UVB IRRADIATION.....	199
7.6	VITAMIN D3 PRODUCTION BY REPEATED LOW-DOSE UVB IRRADIATION	200
7.7	VITAMIN D3 AND UVB-INDUCED SKIN INFLAMMATION	202
7.8	DISCUSSION.....	204
8	CHAPTER 8: FINAL DISCUSSION	213
9	PROSPECTIVE INVESTIGATIONS	222
10	REFERENCES:.....	224

Foreword

Skin cancer is the most common cancer in the world. The majority of skin cancers are caused by exposure of the skin to ultraviolet B radiation (UVB). UVB irradiation causes DNA damage to cells of the skin and suppression of the skin immune system (SIS). UVB irradiation also causes the production of vitamin D3 by skin keratinocytes. Vitamin D3 is enzymatically hydroxylated to produce 25(OH)D3, which is the major circulating form of vitamin D3. 25(OH)D3 is then available for a further hydroxylation to become its biologically active form 1 α ,25 dihydroxyvitamin D3 (1 α ,25(OH)₂D3) by a variety of cells including cells of the skin and the SIS. Keratinocytes of the skin produce 1 α ,25(OH)₂D3 after UVB irradiation and this has led to the discovery that 1 α ,25(OH)₂D3 can suppress the function of key cells of the SIS and protects skin against UVB-induced DNA damage and UVB-induced immunosuppression. This may contribute to a reduced risk of skin cancer development. The focus of this study was to determine whether dietary vitamin D3, and therefore circulating levels of 25(OH)D3 available to be converted to 1 α ,25(OH)₂D3, similarly influenced the development and function of the SIS and the skin's response to UVB-irradiation.

A vitamin D3 deficient mouse population was developed to elucidate the role of dietary vitamin D3 on the development and function of the SIS and response to UVB-irradiation. An immune response known as the contact hypersensitivity (CHS) response, generated by the SIS upon skin exposure to an antigen (contact sensitizer) was compared in vitamin D3 replete and vitamin D3 deficient mice. The investigation of the development and function of the SIS involved an assessment of the directors of the CHS response, dendritic cells (DC); the effectors of the CHS response, T and B cells; and regulators of the CHS, regulatory T cells. UVB-induced suppression of the SIS was also investigated to ascertain the role of dietary vitamin D3 on the skin's response to UVB-irradiation. Vitamin D3 subtly altered DC, CHS effectors and regulatory T cells in an age and gender specific manner. Similarly vitamin D3 protected against UVB-induced immunosuppression in a gender and mouse strain dependent manner. As vitamin D3 deficiency and skin cancer are common, it is important to clarify how vitamin D3 modulates cutaneous biology and how this may relate to skin cancer development.